

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A01N 37/04, 55/02, 59/00, A61K 31/19, 31/28, 33/38	A1	(11) International Publication Number: WO 99/18790 (43) International Publication Date: 22 April 1999 (22.04.99)
(21) International Application Number: PCT/US98/21604 (22) International Filing Date: 9 October 1998 (09.10.98) (30) Priority Data: 60/061,673 10 October 1997 (10.10.97) US (71) Applicant: NVID INTERNATIONAL, INC. [US/US]; 28870 U.S. Highway 19 North, Clearwater, FL 34621 (US). (72) Inventor: ARATA, Andrew, B.; Route 15, Box 1517, Lake City, FL 32054 (US). (74) Agent: FRIJOUF, Robert, F.; Frijouf, Rust & Pyle, P.A., 201 East Davis Boulevard, Tampa, FL 33606 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: DISINFECTANT AND METHOD OF MAKING (57) Abstract A non-toxic environmentally friendly aqueous disinfectant is disclosed for specific use as prevention against contamination by potentially pathogenic bacteria and virus. The aqueous disinfectant is formulated by electrolytically generating silver ions in water in combination with a citric acid. The aqueous disinfectant may include a suitable alcohol and/or a detergent. The aqueous disinfectant has been shown to be very effective at eliminating standard indicator organisms such as staphylococcus aureus, salmonella cholerasuis and pseudomonas aeruginosa.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DISINFECTANT AND METHOD OF MAKING BACKGROUND OF THE INVENTION

Field of the Invention

5 This invention relates to disinfectants and more particularly to an environmentally friendly, non-toxic aqueous disinfectant for specific use against pathogenic bacteria and viruses.

Information Disclosure Statement

10 The prior art has demonstrated that the presence of copper and silver ions in an aqueous solution is useful as a disinfectant. Many in the prior art have used copper and silver ions in an aqueous solution as a disinfectant in water systems such as cooling towers, swimming pools, hot water systems in hospitals, potable water systems, spa pools and the like.

15 Typically, copper and silver electrodes were connected to a direct current power supply. When the direct current was applied to the copper and silver electrodes, copper and silver ions were generated by an electrolysis process from the copper and silver ions within the water. In one example of the prior art, water was passed continuously through an ion chamber having copper and silver electrodes. The water emanating from the ion chamber contained the copper and silver ions generated by copper and silver electrodes within the ion chamber. The water emanating from the ion chamber
20 containing the copper and silver ions was used as a disinfectant in water systems such as cooling towers, swimming pools, hot water systems in hospitals, potable water systems, spa pools and the like. The copper and silver ions within the water systems acted as a disinfectant for controlling algae, viruses, bacteria and the like.

25 U.S. Patent 3,422,183 to Ellison discloses biocide compositions comprising ultra-violet irradiated silver fluoride solutions containing colloidal silver resulting from the irradiation and kept in dispersion by a protective colloid, e.g., casein or gelatin, and biocide uses thereof in slime control, against pathogens or other microbes in food or beverage containers or processing equipment, as an ingredient of wood preservatives, as a bactericide in paints, as a biocide in synthetic polymer films, as a sterilant in
30 bandages, and biocide-like uses in other areas.

U.S. Patent 3,702,298 to Zsoldos discloses a method of maintaining a highly oxidizing aqueous solution intended primarily for treatment of swimming pool water. A metal having a multiple valence is interacted to a lower valence with oxidizable debris in the solution, and the metal is continuously re-oxidized to a higher valence by maintaining in the water a constant excess of an oxidizer bank consisting of a salt of a peroxy acid. Silver, copper and nickel are suitable metals and their salts have germicidal properties which are greatly increased and the spectrum broadened by converting the mono salt to a divalent or trivalent salt.

U.S. Patent 4,180,473 to Maurer et al. discloses a method of transporting metal ions by introducing a metal complex into a medium containing a moiety which demands the metal ion and the complex releases the ions in a controlled manner upon demand. The metal complexes have an aqueous proton induced dissociation property represented by a sigmoidally-shaped curve on a cartesian coordinate plot of the negative log of the metal ion concentration versus the negative log of hydrogen ion concentration. This dissociation property causes a controlled release of metal ion into mediums containing a reacting moiety upon demand for the metal ion. For example, metal working emulsions of oil and water are stabilized by the addition thereto of minor amounts of a metal complex, e.g. disodium monocopper (II) citrate, which at alkaline pH metalworking conditions above about 7 to about 9 releases metal cations to the emulsions imparting stabilizing characteristics which prevent emulsion degradation by a number of factors commonly encountered in metalworking operations. Also, the method is effective in the controlled release of metal ions in the normal range of physiological pH, i.e. about 4 to 9, for growth controlling action against microorganisms including bacteria, fungi and viruses.

U.S. Patent 4,291,125 to Greatbatch discloses a method and apparatus for killing plant and animal bacteria and plant viroids by electrically generated silver ions. The silver ions serve as germicidal agents in infection control and are generated by very slow electrical anodic corrosion of a silver wire located closely adjacent the infection site. In particular, a silver anode and a cathode of non-corroding metal are located in an electrolytic nutrient medium with the silver anode being within five millimeters of the infection site, and a direct voltage is applied to the anode and cathode in a manner passing a positive current in the microampere range into the silver anode causing it

to corrode slightly and give off silver ions which produce a germicidal environment about the infection site.

U.S. Patent 4,385,632 to Odelhog discloses an absorbent body for collecting blood, feces and urine containing a water-soluble copper salt which impedes bacterial growth, prevents the breaking-down of urea into ammonia and complex-binds ammonia
5 so as to prevent the occurrence of unpleasant odor. Preferably copper acetate is used, in which even the acetate ion has germicidal effect.

U.S. Patent 4,564,461 to Skold et al. discloses mechanical working of cast iron performed in the presence of an aqueous metal working composition containing an
10 organic copper (II) complex and an iron corrosion inhibitor. An aqueous concentrate, which after dilution with water is suitable for application in mechanical working of cast iron, contains 1-50% copper (II) complex with such a Cu_2+ content of 0.5-20%, 1-50% iron corrosion inhibitor, 0-50% lubricant, 0-20% pH-regulators, bactericides and solubilizing agents and 10-70% water.

U.S. Patent 4,608,183 to Rossmore discloses antimicrobial mixtures of
15 isothiazolones and a metal complex with a polyfunctional ligand which are synergistic. The mixtures particularly include mixtures of a monocopper disodium citrate as the ligand and a 5-x-2-lower alkyl 4-isothiazolin-3-one wherein x is a halo or hydrogen group as the isothiazolone. The compositions are particularly useful for metal cutting
20 fluids wherein long duration antimicrobial activity is desired.

U.S. Patent 4,666,616 to Rossmore discloses synergistic anti-microbial compositions containing a mixture of a metal complex of a polyfunctional organic liquid and a biocidal composition which contains or releases a lower aldehyde contain-
25 ing 1 to 5 carbon atoms. The compositions are particularly useful as metal working fluids at alkaline pH and have a broad spectrum of activity against fungi and bacterial.

U.S. Patent 4,708,808 to Rossmore discloses synergistic anti-microbial compositions containing a mixture of a metal complex of a polyfunctional organic ligand and a biocidal composition which contains or releases a lower aldehyde contain-
30 ing 1 to 5 carbon atoms. The compositions are particularly useful as metal working fluids at alkaline pH and have a broad spectrum of activity against fungi and bacteria.

U.S. Patent 4,780,216 to Wojtowicz discloses a sanitizing composition consist-
ing essentially of a mixture of a calcium hypochlorite compound and a peroxydisulfate

compound having the formula: $M_xS_2O_8$ where M is an alkali metal or alkaline earth metal, and x is 1 or 2 is employed in treating water to improve pH control and provide increased removal of organic materials. The compositions provide improved sanitation of water in swimming pools, spas, and cooling towers by efficiently oxidizing organic impurities while helping to minimize the increase in the pH of the water. This permits a reduction in the amount and frequency of addition of acidic compounds such as hydrochloric acid to the water bodies. Further, the incorporation of additives such as algaecides, dispersant, and clarifying agents provides for significant improvements in water quality as evidenced by sparkling pure water.

U.S. Patent 4,915,955 to Gomori discloses a concentrate with an unlimited shelf-life, which can be mixed with hydrogen peroxide at a ratio of 1:99 to 1:199 to become an effective disinfectant, is obtained when a viscous solution of inorganic acid, with a pH less than or equal to 1.6, is mixed with a silver salt compound or a colloidal silver compound at 50° to 66° C. The mixture is further combined at room temperature with other inorganic acid(s) to reach a total of 100 g inorganic acid(s) per liter of water at room temperature, an organic acid stabilizer is added and the mixture is homogenized. The concentrate, during storage, remains homogeneous and crystal-clear.

U.S. Patent 4,933,178 to Capelli discloses a medical device with an antimicrobial coating that is safe, effective, photostable and readily manufacturable produced by applying a composition to at least one body fluid-contacting surface of the device such that a solid coating is provided on that surface, the coating composition comprising an oligodynamic metal salt of a sulfonylurea, a polymeric material, at least one acid compound selected from the group consisting of a water-soluble carboxylic acid and water-insoluble carboxylic acid, and a carrier liquid in which foregoing components are soluble. The antimicrobial coating accommodates variation in the release of antimicrobial metal ions as a function of the intended use for a medical device to which the coating is applied.

U.S. Patent 5,017,295 to Antelman discloses a method or methods of controlling the growth of bacteria in the water of swimming pools and/or industrial water supplies by adding to the water a specified concentration of a stable divalent silver compound. The invention has the advantage over chlorination in that it is odorless and non-volatile. It furthermore is superior to monovalent silver compounds as these compounds do not

decompose in the presence of light and resist precipitation by halides and form divalent soluble complexes which in the monovalent state are invariably insoluble solids.

U.S. Patent 5,073,382 to Antelman discloses a solid alkaline bactericidal compositions suitable for compounding alkaline end products such as food and dairy
5 cleaners and surgical scrubbing soaps, formed by the neutralization of acid stabilized inorganic divalent silver complexes and capable of effecting 100% kills upon cultures of anaerobic bacteria colonies of 100K/cc. within 5 minutes.

U.S. Patent 5,078,902 to Antelman discloses divalent silver halides providing a source for divalent bactericidal silver ions in the presence of persulfate. The halides
10 are especially effective when applied to water used in industrial cooling installations, hot tubs and swimming pools and will conform to stringent EPA requirements for waters utilized for bathing as in tubs and pools of 100% kills of 100 K/cc E. Coli coliforms within 10 minutes, exemplary of which are the chloride and bromide which give 100% kills within 5 minutes. The halides, of course, can be used in salty water
15 since they are solids immune from halide action that would otherwise precipitate soluble divalent silver from solution.

U.S. Patent 5,089,275 discloses solid bactericidal compositions based on divalent silver (Ag(II)) as the active sanitized agent. The compositions are prepared by reacting acid liquid Ag(II) complexes with anhydrous calcium sulfate so as to form
20 a solid matrix in which the bactericide is entrapped in the resulting hydrated calcium sulfate. Optimum compositions are described consisting of Ag(II) of solid (by weight) to liquid (by volume) is 5:2. The resulting solid bactericides can be used in water cooling installations. They are capable of causing 100% kills within 10 minutes of E. Coli conforms in conformity with EPA protocols, allowing them to qualify as
25 swimming pool and hot tub sanitizers. Since the compositions are based on calcium sulfate, they are also suitable as mineralizers, thus providing a dual function.

U.S. Patent 5,332,511 to Gay et al. discloses a process for sanitizing water in swimming pools, spas and hot tubs whereby the level of bacteria in said water is lowered comprising treating said water with a bactericidal effective amount of a
30 combination of diisodecyl dimethyl ammonium chloride and copper (II) ions, the concentration of diisodecyl dimethyl ammonium chloride in said water being less than about 60 parts per million parts of water by weight and treating said water at least

intermittently with an oxidant selected from the group consisting of available chlorine and ozone.

U.S. Patent 5,364,649 to Rossmore et al. discloses activity of antimicrobial compounds selected from isothiazolones and compounds which release formaldehyde enhanced with a metal complex of a lower alkanolamine, particularly copper (cupric) trietha-iolamine. The enhancement is particularly useful in metalworking fluids.

U.S. Patent 5,373,025 to Gay discloses a sanitizer composition comprising a bactericidal effective amount of the combination of (a) a quaternary ammonium compound selected from the group consisting of (hydrogenated tallow) 2-ethylhexyl dimethyl ammonium salt, dicoco dimethyl ammonium salt, and mixtures thereof; and (b) a copper (II) ion source.

U.S. Patent 5,382,337 to Wlassics et al. discloses a process for oxidizing organic materials or compounds in aqueous phase, with hydrogen peroxide and in the presence of ferrous ions $FE-(II)$, and optionally cupric ions $cu-(II)$, carried out under irradiation with artificial visible light.

U.S. Patent 5,464,559 to Marchin et al. discloses a composition provided for treating drinking water for disinfecting and/or removing iodide. The composition utilizes resin bound silver ions. For performing the disinfection or iodide removal with minimal release of silver ions into the water being treated, a chelating resin having iminodiacetate chelating groups is employed, and the resin is loaded with not over 0.5 mole of silver ions per mole of iminodiacetate.

U.S. Patent 5,503,840 to Jacobson et al. discloses an antimicrobial composition of titanium dioxide, barium sulfate, zinc oxide particles, and mixtures thereof having successive coatings of silver, in some cases a coating of zinc and/or copper compounds such as zinc oxide, copper (II) oxide and zinc silicate; silicon dioxide; alumina; and a dispersion aid such as dioctyl azelate.

U.S. Patent 5,510,109 to Tomioka et al discloses an antibacterial and antifungal composition which comprises an antibacterial and antifungal material carried on a porous particle carrier. Preferably, the porous particle carrier is a silica gel particle. The antibacterial and antifungal material is at least one metal complex salt, and can contain plant extracts and the like in addition to the metal complex salt. At least a

portion of the surface of the above-mentioned carrier having the antibacterial and antifungal composition can be coated with a coating material.

Unfortunately, these copper and silver ions within an aqueous solution have only a limited stable ionic life. After a limited time, the copper and silver ions form complexes with other elements thus diminishing the concentration of the copper and silver ions within the aqueous solution. Accordingly, the aqueous solution had to be replenished with copper and silver ions to maintain the concentration of the copper and silver ions within the aqueous solution. The aqueous solution may be replenished with copper and silver ions by constantly circulating the aqueous solution thorough the ion chamber.

The present invention provides an aqueous disinfectant solution having a stable ionic form having an extended useful shelf-life. The extended useful shelf-life of the aqueous disinfectant solution enables the aqueous disinfectant solution to be packaged in an aqueous concentrate form.

Therefore, it is an object of the present invention to provide an improved disinfectant and the method of making comprising an aqueous disinfectant for specific use as prevention against contamination by potentially pathogenic bacteria and virus and antifungal properties.

Another object of this invention is to provide an improved disinfectant and the method of making which is an effective disinfectant for eliminating standard indicator organisms such as staphylococcus aureus, salmonella cholerasuis and pseudomonas aeruginosa.

Another object of this invention is to provide an improved disinfectant and the method of making which is a non-toxic, environmentally friendly aqueous disinfectant.

Another object of this invention is to provide an improved disinfectant and the method of making which comprises a stable ionic formulation having an extended useful shelf-life.

Another object of this invention is to provide an improved disinfectant and the method of making which may be packaged in a concentrated aqueous form.

Another object of this invention is to provide an improved disinfectant and the method of making which may be electrolytically generated in a batch process or a continuous process.

Another object of this invention is to provide an improved disinfectant and the method of making which is electrolytically generated in an economical manner.

Another object of this invention is to provide an improved disinfectant and the method of making which is suitable for use with an alcohol and/or a detergent.

5 Another object of this invention is to provide an improved disinfectant and the method of making which may be used on exposed and/or contaminated surfaces to kill bacteria, virus, fungi and other micro-organisms.

10 Another object of this invention is to provide an improved disinfectant and the method of making which may be used on contaminated open wounds and tissue, dermal wound sites and/or lesions of living organisms such as animals and humans.

Another object of this invention is to provide an improved disinfectant and the method of making which may be used on exposed surfaces in food processing plants, residential, hospital, restaurants, public facilities and the like.

15 The foregoing has outlined some of the more pertinent objects of the present invention. These objects should be construed as being merely illustrative of some of the more prominent features and applications of the invention. Many other beneficial results can be obtained by applying the disclosed invention in a different manner or modifying the invention within the scope of the invention. Accordingly other objects in a full understanding of the invention may be had by referring to the summary of
20 the invention, the detailed description describing the preferred embodiment in addition to the scope of the invention defined by the claims taken in conjunction with the accompanying drawings.

SUMMARY OF THE INVENTION

A specific embodiment of the present invention is described and shown in the attached Detailed Description. For the purpose of summarizing the invention, the invention relates to an improved non-toxic environmentally friendly aqueous disinfectant for use as a prevention against contamination by potentially pathogenic bacteria, virus and fungi. The improved aqueous disinfectant is suitable for use on exposed surfaces. In addition, the improved aqueous disinfectant is suitable for use on dermal wound sites and lesions of living organisms such as animals and humans. The aqueous disinfectant is pH neutral.

The improved aqueous disinfectant comprises an aqueous solution of silver citrate wherein the silver is electrolytically generated in a solution of citric acid and water. The electrolytically generated silver forms an organic metal complex with the citric acid such as a chelated organic metal complex with the citric acid. In one example of the invention, the solution of citric acid and water comprises approximately 5.0% to 10.0% citric acid by volume. The silver citrate formed by the electrolytically generated silver has a concentration in excess of 0.0005% by volume.

In another example of the invention, the invention is incorporated into an aqueous disinfectant in a concentrated form having an extended shelf-life comprising an aqueous solution of silver citrate wherein the silver is electrolytically generated in a solution of citric acid in water. The electrolytically generated silver has a concentration of in excess of 0.05% by volume.

The aqueous disinfectant may be combined with an alcohol such as ethyl alcohol (ETOH) and/or a detergent such as sodium dodecyl sulfate.

The invention is also incorporated into the process of making the disinfectant comprising the step of electrolytically generating silver in a solution of citric acid and water to form an aqueous solution of silver citrate. The process may include creating a solution of approximately 5.0% to 10% citric acid in water by volume. A positive silver electrode is spaced relative to a negative electrode for enabling the solution to be located therebetween. A potential difference is applied to the positive and negative electrodes to establish a flow of silver ions between the positive and negative electrodes for enabling the silver ions to react with the citric acid to form silver citrate thereby.

The invention is also incorporated into the process of making silver citrate, comprising the step of electrolytically generating silver in a solution of citric acid and water to form an aqueous solution of silver citrate.

5 The foregoing has outlined rather broadly the more pertinent and important features of the present invention in order that the detailed description that follows may be better understood so that the present contribution to the art can be more fully appreciated. Additional features of the invention will be described hereinafter which form the subject of the invention. It should be appreciated by those skilled in the art that the conception and the specific embodiments disclosed may be readily utilized as
10 a basis for modifying or designing other structures for carrying out the same purposes of the present invention. It should also be realized by those skilled in the art that such equivalent constructions do not depart from the spirit and scope of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

For a fuller understanding of the nature and objects of the invention, reference should be made to the following detailed description taken in connection with the accompanying drawings in which:

5 FIG. 1 is a diagram of a first process of making the disinfectant of the present invention;

FIG. 2 is a diagram of a second process of making the disinfectant of the present invention;

FIG. 3 is an enlarged detailed view of the ion chamber of FIGS. 1 and 2;

10 FIG. 4 is an enlarged detailed view of an ion chamber suitable for making the disinfectant of the present invention in a batch process;

FIG. 5 is a table illustrating the shelf-life tests for initial sampling intervals;

FIG. 6 is a table illustrating the shelf-life tests for secondary sampling intervals;

FIG. 7 is a table illustrating the efficacy tests against salmonella cholerasuis;

15 FIG. 8 is a table illustrating the efficacy tests against staphylococcus aureus;
and

FIG. 9 is a table illustrating the efficacy tests against pseudomonas aeruginosa.

Similar reference characters refer to similar parts throughout the several Figures of the drawings.

DETAILED DISCUSSION

PROCESS OF MAKING

FIG. 1 is a diagram of a first process 10 of making the disinfectant 14 of the present invention. The first process 10 is shown as a continuous process of making the disinfectant 14. It should be understood that the first process 10 of FIG. 1 is only an example of a process and numerous other variations and/or processes may be utilized to make the disinfectant 14 of the present invention.

The disinfectant 14 may be used immediately for any suitable application such as a disinfectant in a water system including cooling towers, hot water systems, potable water systems, or any other suitable application or surface.

The first process 10 comprises a water input conduit 16 for introducing water 18 from a water source (not shown) to a water treatment unit shown as a reverse osmosis unit 20. The reverse osmosis unit 20 passes the water 18 from the water input conduit 16 through a semi-permeable membrane (not shown) for removing impurities from the water. Although the water treatment unit is shown as a reverse osmosis unit 20 it should be understood that various water treatment units may be employed within the process shown in FIG. 1. Preferably, the water 18 emanating from the reverse osmosis unit 20 is deionized medically pure water.

The water 18 emanating from the reverse osmosis unit 20 is directed to a valve 30 through a conduit 31. The valve 30 directs the water 18 through a conduit 32 to a flow control injector 40. A citric acid tank 50 contains concentrated citric acid. The concentrated citric acid is directed by a conduit 51 to a metering valve 60 for metering the concentrated citric acid into the flow control injector 40. The flow control injector 40 mixes the concentrated citric acid with the water 18 to provide a dilute citric acid solution 62. The metering valve 60 controls the concentration of the citric acid within the water 18. The diluted citric acid solution 62 is directed by a conduit 62 into an ion chamber 70.

FIG. 3 is an enlarged detailed view of the ion chamber 70 of FIG. 1. The ion chamber 70 includes a positive and a negative electrode 71 and 72. The positive and negative electrodes 71 and 72 are located in a spaced apart position for enabling the diluted citric acid solution 62 to pass between the positive and negative electrodes 71

and 72. Each of the positive and negative electrodes 71 and 72 is fabricated from elemental silver. Preferably, the positive and negative electrodes 71 and 72 are formed from 99.9999% pure elemental silver.

5 A direct current power supply 80 includes a positive and a negative conductor 81 and 82 connected to the positive and negative electrodes 71 and 72. The positive and negative electrodes 71 and 72 are spaced apart a suitable distance such as 2.0 to 8.0 centimeters to allow an ionic current flow between the positive and negative electrodes 71 and 72.

10 Upon energizing the direct current power supply 80, an ion current flows between the positive and negative electrodes 71 and 72. The direct ion current flow between the positive and negative electrodes 71 and 72 produces electrolytically free silver ions within the diluted citric acid solution 62. The silver ions react with the citric acid in the diluted citric acid solution 62 to produce the disinfectant 14 of the present invention.

15 The disinfectant 14 is directed by a conduit 86 to a settling tank 90. The settling tank 90 includes an overflow conduit 91 and a drain conduit 92. The disinfectant 14 exits the settling tank 90 through the overflow conduit 91. Any precipitated materials from the disinfectant 14 within the settling tank 90 fall to the bottom of the settling tank 90. The precipitated materials at the bottom of the settling tank 90 may
20 be removed through the drain conduit 92 to a purge tank 100. The precipitated materials in the purge tank 100 may be recycled.

The disinfectant 14 exiting through the overflow conduit 91 from the settling tank 90 is directed to a particle filter 110. Although the particle filter 110 may be any suitable filter, preferably the particle filter 110 is a submicron filter. The filtered
25 disinfectant 14 is directed to a valve 120 by a conduit 121. The valve 120 directs the filtered disinfectant 14 to a conduit 122 for discharge from the first process 10.

The filtered disinfectant 14 discharged from conduit 122 may be used immediately for any suitable application such as a disinfectant in a water system or any other suitable application. In the event a greater concentration of the disinfectant 14 is
30 desired, the disinfectant 14 may be recirculated for increasing the concentration of the disinfectant 14.

FIG. 2 is a diagram of a second process 10A of making the disinfectant 14 of the present in a concentrated form. The second process 10A is shown as a recirculating process of making the disinfectant 14 and for increasing the concentration of the disinfectant 14. In the concentrated form, the disinfectant 14 may be bottled for use at a later time. It should be understood that the second process 10A of FIG. 2 is only an example of a process and numerous other variations and/or processes may be utilized to make the disinfectant 14 of the present invention.

In the second process 10A shown in FIG. 2, the valve 30 and 120 are move into positions opposite to the positions shown in FIG. 1. The valve 120 directs the filtered disinfectant 14 to a conduit 123. The conduit 123 is connected through a conduit 130 to the conduit 32 of the valve 30.

The valve 30 directs the filtered disinfectant 14 though the conduit 32 to the flow control injector 40. Additional concentrated citric acid is directed through the metering valve 60 into the flow control injector 40. The flow control injector 40 mixes the concentrated citric acid with the filtered disinfectant 14 to increase the concentration of the citric acid solution 62A.

The citric acid solution 62A is directed into an ion chamber 70 to produce additional silver ions within the citric acid solution 62A. The silver ions react with the citric acid in the citric acid solution 62A to increase the concentration of the disinfectant 14. The disinfectant 14 is passed through the settling tank 90 to exit through the overflow conduit 91. The disinfectant 14 is filtered by the particle filter 110 and is directed to the valve 120 by the conduit 121.

The valve 30 and 120 are maintained in positions shown in FIG. 2 to continue to recirculate the disinfectant 14 for increasing the concentration of the disinfectant 14. Upon obtaining the desired concentration of the disinfectant 14, the valve 120 may be moved to the position shown in FIG. 1 to discharge the disinfectant 14 from the conduit 122.

FIG. 4 is an enlarged detailed view of an ion chamber 170 suitable for making the disinfectant of the present invention in a batch process. The ion chamber 170 includes a positive and a negative electrode 171 and 172. Each of the positive and negative electrodes 171 and 172 is fabricated from 99.9999% pure elemental silver.

The positive and negative electrodes 171 and 172 are located in a spaced apart position for enabling the citric acid solution 162 to pass between the positive and negative electrodes 171 and 172. Preferably, the positive silver electrode 171 is spaced relative to a negative electrode 172 a distance sufficient to enable silver ion flow therebetween. The spacing of the positive and negative electrodes 171 and 172 has been shown in an exaggerated fashion in FIG. 4. Preferably, a spacing of approximately 2.0 to 8.0 mm. has been found to be suitable for the above concentration of citric acid and water.

A direct current power supply 180 includes a positive and a negative conductor 181 and 182 connected to the positive and negative electrodes 171 and 172. Upon energizing the direct current power supply 180, an ion current flows between the positive and negative electrodes 171 and 172. The direct ion current flow between the positive and negative electrodes 171 and 172 produces electrolytically free silver ions within the citric acid solution 162. The silver ions react with the citric acid in the citric acid solution 162 to produce the disinfectant 14 of the present invention.

The process of making a disinfectant comprises electrolytically generating silver ions in a solution of citric acid and water to form an aqueous solution of silver citrate. Preferably, the solution of citric acid and water comprises a solution of approximately 5.0% to 10% citric acid in water by volume. A potential difference of 12 volts to 50 volts provides a flow of silver ions in the range of 0.1 amperes to 0.5 amperes per square inch. A more fuller explanation of the content of the solution within the ion chamber 170 will be described in greater detail hereinafter.

The prior art has established in that the generation of both silver ions and copper ion in water provides the best disinfectant properties. The combination of silver ions and copper ions provides superior disinfecting properties than either silver ions alone or copper ions alone. This synergistic effect of silver ions and copper ions in water has been well established by the prior art.

In contrast to this established prior art, the disinfectant of the present invention is formed in a solution of citric acid and water rather than water alone. Additionally, the disinfectant of the present invention has superior properties with only silver ions alone rather than the combination of both silver ions and copper ions. The silver ions of the present process react with the citric acid to form a silver citrate. The silver

citrate provides superior disinfectant properties over the prior art process of generating silver and copper ions in water.

In further contrast to the established prior art, the disinfectant of the present invention has a stable ionic form having an extended useful shelf-life. The useable shelf-life of the disinfectant of the present invention enables the aqueous disinfectant solution to be packaged in an aqueous concentrate form.

COMPOSITION

The improved disinfectant is an aqueous solution of silver citrate wherein the silver is electrolytically generated in a solution of citric acid and water. The silver citrate formed in accordance with the above process has different characteristics than other forms of silver citrate.

Concentrations of 0.1% silver citrate by volume have been formulated in accordance with the above process. A concentration of 0.1% silver citrate by volume corresponds to 1000 parts per million (ppm). The concentration of 0.1% silver citrate was formed in a solution of citric acid and water comprises approximately 10.0% citric acid by volume. Higher concentration of the silver citrate are believed to be obtainable by the above process. It appears the higher the concentration of citric acid in water, the higher the concentration of silver citrate formed by the above process.

The Merck Index, Eleventh Edition (1989) page 1348 states that silver citrate is soluble in 3500 parts water. A concentration of 1 to 3500 corresponds to 285 parts per million (ppm). Obviously, the silver citrate formed in accordance with the above process has different solubility than other forms of silver citrate.

Nuclear magnetic resonance tests (1H NMR) were performed on the silver citrate formed in accordance with the above process and a blank citric acid sample. The samples showed an overwhelming excess of citric acid, with little or no other anions present. It is postulated the Ag must be in the form of the cation Ag⁺ complexed with the citric acid. It is theorized the empty 5s orbital of Ag⁺ overlaps with the delocalized π bond on one of the carboxyl groups of citric acid. The citric acid anion is the counterion for this complex ion (Ag(CA)_x)⁺ i.e. (CA). CA is citric acid or is (C₆H₈O₇ - H₂O). Another possibility is a zwitterion, where the negative charge is on the complex itself, (Ag+CA⁻) where the total charge of the complex is neutral. Either

or both of these species may exist in the silver citrate formed in accordance with the above process. Multiple complexation to Ag^+ is also possible.

A second formulation of the improved disinfectant of the present invention includes the addition of an alcohol. In one example of the second formulation of the improved disinfectant, ethyl alcohol (ETOH) is added in an approximate amount of 20% by volume. However, it should be understood that other types of alcohols may be added to the second formulation of the improved disinfectant of the present invention.

A third formulation of the improved disinfectant of the present invention includes the addition of a detergent. In one example of the third formulation of the improved disinfectant, sodium dodecyl sulfate is added in an approximate amount of 0.1% by volume.

SHELF-LIFE STUDY

The copper and silver ions in the prior art aqueous solution have only a limited stable ionic life. After a limited time, the copper and silver ions in the prior art aqueous solution form complexes with other elements thus diminishing the concentration of the copper and silver ions within the aqueous solution.

A significant difference of the disinfectant of the present invention is the stable life of the silver citrate. The present invention provides an aqueous disinfectant solution having a stable ionic form having an extended useful shelf-life. The extended useful shelf-life of the disinfectant of the present invention enables the disinfectant to be packaged in an aqueous concentrate form.

A series of tests was preformed on the following formulations.

1. Silver and Citric Acid (1.0% citric acid solution/pH 6.0)
2. Silver and Citric Acid (5.0% citric acid solution/pH 6.0)
3. Silver and Citric Acid (10% citric acid solution/pH 6.0)

The silver and citric acid formulations were prepared using 100/100 silver:silver electrodes. The electrodes were immersed in 1.0, 5.0 and 10% citric acid solutions and a current was applied for approximately two hours. The solutions were stored for 24 hours to allow for precipitation. The solutions were filtered using #2 Whatman

filter paper. The final pH was adjusted to 6.0 with sodium carbonate and sodium bicarbonate.

FIG. 5 is a table illustrating the results of the shelf-life test for the initial shelf-life sampling intervals. The initial intervals for the initial shelf-life sampling intervals of the disinfectant were 1 week, 2 weeks, 3 weeks and 4 weeks. FIG. 5 illustrates that silver citrate is not stable at high concentrations in the 1.0% citric acid solution. The 300 ppm silver citrate did not remain in the 1.0% citric acid solution. However, the 300 ppm silver citrate was stable in the 10% citric acid solution.

FIG. 6 is a table illustrating the results of the shelf-life test for secondary shelf-life sampling intervals. The secondary intervals for the secondary shelf-life sampling intervals of the disinfectant were 0 weeks, 7 weeks, 14 weeks and 21 weeks. FIG. 6 also illustrates that silver citrate is not stable at high concentrations in the 1.0% citric acid solution. Conversely, the silver citrate was stable in both the 5% and 10% citric acid solutions.

The results seen in FIG. 6 for week 21 confirm the stability of the silver citrate in the 5.0% and 10% citric acid solutions. The stability of the silver citrate in the 1.0% citric acid solution experienced significant reductions during the last phase of the study. The minimum concentration of the citric acid solution is therefore some value greater than 1.0% and less than 5.0%. The maximum concentration of the citric acid in the aqueous solution has not been determined by test. However, it is believed that the maximum concentration of the citric acid in the aqueous solution much greater than 10.0%. It is also evident from these results, that the higher the concentration of the citric acid in the aqueous solution, the greater the concentration of silver ions that can be stabilized.

LABORATORY STUDY

In order to establish the effectiveness of the improved disinfectant of the present invention, laboratory tests were performed against various test microorganisms. The test microorganisms considered were (a) *Pseudomonas aeruginosa* strain ATCC 15442, (b) *Salmonella choleraesuis* strain ATCC 10708 and (c) *Staphylococcus aureus* strain ATCC 6538.

The inoculum level for each of the test microorganisms were established in a similar manner. Test strains were grown individually at 35° C for 24 hr. The cells were harvested by centrifugation at 10,000 x g for 10 minutes and washed twice with Butterfield's Phosphate Buffer (BPB of pH 7.2). The cells were resuspended in the
5 Butterfield's Phosphate Buffer to obtain a cell suspension of approximately 1.0×10^8 CFU/mL for each microorganism (target inoculum levels were approx. 10^6 in the final test solution).

The test microorganisms considered were tested at uniform sampling intervals. The sampling intervals selected were (a) 15 seconds (ethanol trials only), (b) 1 minute,
10 (c) 5 minutes, (d) 10 minutes and (e) 30 minutes.

Five compounds were tested against the test microorganisms. The five compounds tested were (a) silver and citric acid (4.27 ppm in a 0.1% citric acid solution), (b) copper and citric acid (4.07 ppm in a 0.1% citric acid solution), (c) citric acid (0.1% citric acid solution), (d) silver (4.08 ppm), citric acid (0.1%) and ethanol (20%)
15 and (e) Ethanol (20%).

The silver and citric acid (4.27 ppm in a 0.1% citric acid solution) was prepared using 100/100 silver:silver electrodes. The electrodes were immersed in a 0.1% citric acid solution and current was applied for approximately two hours. The solution was stored for 24 hours to allow for precipitation. The solution was filtered using No. 2
20 Whatman filter paper. The final pH was adjusted to 7.0. The concentration tested had a silver concentration of 4.27 mg/L.

The copper and citric acid (4.07 ppm in a 0.1% citric acid solution) was prepared using 100/100 copper:copper electrodes. The electrodes were immersed in a 0.1% citric acid solution and current was applied for approximately two hours. The
25 solution was stored for 24 hours to allow for precipitation. The solution was filtered using #2 Whatman Filter paper. The final pH was adjusted to 7.0. The concentration tested had a copper concentration of 4.07 mg/L (as measured by ICAP).

The citric acid (0.1% citric acid solution) was prepared using deionized water. The pH was adjusted to 7.0.

30 The silver (4.08 ppm), citric Acid (0.1%) and ethanol (20%) was prepared using 100/100 silver:silver electrodes. The electrodes were immersed in a 0.1% citric acid solution and current was applied for approximately two hours. The solution was stored

for 24 hours to allow for precipitation. The solution was filtered using #2 Whatman filter paper. The final pH was adjusted to 7.0. The solution was diluted with ethanol to yield a concentration of 4.08 mg/L silver in a 20% ethanol solution.

5 The Ethanol (20%) was prepared with by diluting Reagent grade ethanol with deionized water to make the appropriate dilution.

The test microorganisms were tested in accordance with the following test procedures. Duplicate trials were conducted for each test variable. Ninety nine volumes of the test solutions in 250 mL Erlenmeyer flasks were prepared from sterilized deionized water. The solutions were inoculated separately with one mL of
10 24 hour culture from each of the test strains to yield a flask inoculum level of approximately 1.0×10^6 CFU/mL. The actual count for each of the microorganisms are set forth in FIGS. 7-9.

Solutions were mixed well and kept under constant agitation. Samples of 1.0 mL were removed at the above specified time intervals and placed into 9.0 mL
15 Neutralization Broth media (Difco) to yield an initial dilution of 1:10. All samples were serially diluted in the Butterfield's Phosphate Buffer solution (BPB) and plated onto Tryptic Soy Agar (TSA) in duplicate using the pour plate technique. Percent reductions were calculated for each test solution against each test strain.

The results of the laboratory study can be seen in FIGS. 7-9. For all tests
20 which utilized either copper or silver ions, concentrated solutions were prepared 24 hours prior to the beginning of the study. Solutions were filtered and determinations for ion content were made. From these stock solutions (copper ion concentration as measured by ICAP and silver ion concentration as measured by Atomic Absorption analysis), final working solutions were made. The target ion concentration for both
25 copper and silver was 5.0 mg/L.

FIG. 7 is a table illustrating the efficacy tests against salmonella cholerasuis. The trials that utilized 20% ethanol showed a slow, but complete disinfection. The ethanol solution has an approximate $1.0 \log_{10}$ reduction after one minute. Near complete disinfection was seen after 30 minutes of contact time. Of the three organisms tested, salmonella cholerasuis was the one most effected by the ethanol disinfectant. The copper: citric acid was not effective in disinfecting salmonella cholerasuis at any of time periods. The citric acid solution was slightly more effective in reducing
30

the number of salmonella cholerasuis, achieving a 1.0 log₁₀ reduction at the 30 minute time period. Both silver: citric acid and silver: citric acid with ethanol exhibited a 6.0 log₁₀ reduction over the course of the 30 minute trial. The silver: citric acid solution showed a 5.0 log₁₀ reduction within the first 5 minutes and a greater 6.0 log₁₀ reduction at the 10 minute time period. Silver: citric acid with ethanol appeared to be the most effective, exhibiting a 2.36 log₁₀ reduction within in the first minute and a greater than 6.0 log₁₀ reduction within the first 5 minutes of contact.

FIG. 8 is a table illustrating the efficacy tests against staphylococcus aureus. This table indicates a different reaction for the 20% ethanol against staphylococcus aureus as compared to salmonella cholerasuis. No significant reduction was seen between 15 seconds and 30 minutes. Neither citric acid nor copper: citric acid was effective against staphylococcus aureus. Neither of the aforementioned formulas were able to significantly reduce the number of staphylococcus aureus organisms present within the 30 minute time period. However, both silver: citric acid and silver: citric acid with ethanol exhibited a 6.0 log₁₀ reduction over the course of the 30 minute trial. The silver: citric acid solution showed a 3.0 log₁₀ reduction within the first 10 minutes and a greater than 6.0 log₁₀ reduction at the end of 30 minutes. Silver: citric acid with ethanol appeared to be the most effective, exhibiting a 2.36 log₁₀ reduction within the first minute and a greater than 6.0 log₁₀ reduction within the first 5 minutes of contact.

FIG. 9 is a table illustrating the efficacy tests against pseudomonas aeruginosa. The seen in this table for pseudomonas aeruginosa, indicate similar results as those seen for that used staphylococcus aureus. For the 20% ethanol trials, no significant reduction was seen between 15 seconds and 30 minutes. This same trend was recorded for citric acid and copper: citric acid. Both silver: citric acid and silver: citric acid with ethanol exhibited near or greater than 6.0 log₁₀ reductions over the course of the 30 minute trial. The silver: citric acid solution showed a 2.49 log₁₀ reduction at the 10 minute time period and a greater than 5.70 log₁₀ reduction at the end of 30 minutes. Silver citric acid with ethanol showed the best disinfection against pseudomonas aeruginosa, mirroring the results seen with the other two organisms. A greater than 6.0 log₁₀ reduction was recorded at the 5 minute sampling period.

FIELD TRIAL RESULTS

5 The improved disinfectant has been tested in preliminary veterinary field trials to establish the effectiveness of the present invention. The veterinary field trial test were conducted by licensed veterinarians on equine species. The improved disinfectant was tested on contaminated open, non-healing tissue and wounds. The open, non-healing wounds were treated with wet dressings or by spraying the improved disinfectant onto the wound.

10 The disinfectant has been tested on dermal lesions both contaminated and infected with gram negative and gram positive bacteria. The results have shown that this formulation exhibits superior performance as compared to available disinfectant products currently on the market. The disinfectant formulation has shown to be very efficacious for irrigating deep wounds and abscesses without damage to tissue. Decreased healing time and reduction in scar formation have been observed repeatedly during the study. The disinfectant appears to promote healthy granulation without
15 excessive fibrosis.

The disinfectant has been used as a surface disinfectant and therefore has shown best results with extended contact with the contaminated tissue. On surface wounds, best results are obtained with "wet dressing" or frequent spray applications for dermal surfaces not amenable to applied dressing. Drained abscesses are flushed, the disinfectant solution is held in the cyst, then drained and again filled and agitated for 2-3
20 minutes before allowing to drain. Deep wounds closed with drains have shown rapid healing time and reduced draining when flushed with the disinfectant. An additional use for the disinfectant may be as a uterine flush for bacterial and/or fungal/yeast infection. Preliminary results with this application have shown to be very promising.

25 The present disclosure includes that contained in the appended claims as well as that of the foregoing description. Although this invention has been described in its preferred form with a certain degree of particularity, it is understood that the present disclosure of the preferred form has been made only by way of example and that numerous changes in the details of construction and the combination and arrangement
30 of parts may be resorted to without departing from the spirit and scope of the invention.

WHAT IS CLAIMED IS:

1. An aqueous disinfectant, comprising:
an aqueous solution of silver citrate wherein the silver is electrolytically generated in a solution of citric acid and water.
2. An aqueous disinfectant as set forth in claim 1, wherein the electrolytically generated silver forms an organic metal complex with the citric acid.
3. An aqueous disinfectant as set forth in claim 1, wherein the electrolytically generated silver forms a chelated organic metal complex with the citric acid.
4. An aqueous disinfectant as set forth in claim 1, wherein the electrolytically generated silver forms a complex with the citric acid of $(\text{Ag}(\text{CA})_x)^+ (\text{CA})^-$, wherein CA is $(\text{C}_6\text{H}_8\text{O}_7 - \text{H}_2\text{O})$.
5. An aqueous disinfectant as set forth in claim 1, wherein the electrolytically generated silver forms a complex with the citric acid of $(\text{Ag}+\text{CA})^-$, wherein CA is $(\text{C}_6\text{H}_8\text{O}_7 - \text{H}_2\text{O})$.
6. An aqueous disinfectant as set forth in claim 1, wherein the solution of citric acid and water comprises approximately 5.0% to 10.0% citric acid by volume.
7. An aqueous disinfectant as set forth in claim 1, wherein the solution of citric acid and water comprises approximately 5.0% to 10.0% citric acid by volume; and
approximately 0.0005% to 0.001% by volume of silver citrate formed by the electrolytically generated silver.
8. An aqueous disinfectant in concentrated form having an extended shelf-life, comprising:

an aqueous solution of silver citrate wherein the silver is electrolytically generated in a solution of citric acid in water; and

the electrolytically generated silver having a concentration of in excess of 0.05% by volume.

9. An aqueous disinfectant in concentrated form having an extended shelf-life, comprising:

an aqueous solution of silver citrate wherein the silver is electrolytically generated in a solution of approximately 5.0% to 10.0% by volume of citric acid in water; and

the electrolytically generated silver having a concentration of approximately 0.05% to 0.1% by volume.

10. An aqueous disinfectant, comprising:

an aqueous solution of silver citrate in a solution of citric acid and water wherein the concentration of silver citrate exceeds 0.05% by volume.

11. An aqueous disinfectant, comprising:

an aqueous solution of silver citrate formed from electrolytically generated silver in a solution of citric acid and water; and

approximately 20% alcohol by volume.

12. An aqueous disinfectant as set forth in claim 11, wherein the alcohol is approximately 20% ethyl alcohol (ETOH) by volume.

13. An aqueous disinfectant as set forth in claim 11, wherein the solution of citric acid and water comprises approximately 5.0% to 10.0% citric acid by volume.

14. An aqueous disinfectant as set forth in claim 11, wherein the solution of citric acid and water comprises approximately 5.0% to 10.0% citric acid by volume; and

the electrolytically generated silver comprising approximately 0.0005% to 0.001% by volume.

15. An aqueous disinfectant as set forth in claim 11, wherein the solution of citric acid and water comprises approximately 5.0% to 10.0% citric acid by volume; and
the electrolytically generated silver comprising approximately 0.05% to 0.1% by volume.
16. An aqueous disinfectant, comprising:
an aqueous solution of silver citrate formed from electrolytically generated silver in a solution of citric acid and water;
approximately 20% ethyl alcohol by volume; and
approximately 0.01% to 0.1% anionic detergent by volume.
17. An aqueous disinfectant as set forth in claim 16, wherein the solution of citric acid and water comprises approximately 5.0% to 10.0% citric acid by volume.
18. An aqueous disinfectant as set forth in claim 16, wherein the solution of citric acid and water comprises approximately 5.0% to 10.0% citric acid by volume; and
approximately 0.0005% to 0.001% by volume of silver citrate formed by the electrolytically generated silver.
19. An aqueous disinfectant as set forth in claim 16, wherein the solution of citric acid and water comprises approximately 5.0% to 10.0% citric acid by volume; and
the electrolytically generated silver comprising approximately 0.05% to 0.1% by volume.
20. An aqueous disinfectant as set forth in claim 16, wherein the detergent is sodium dodecyl sulfate.
21. The process of making a disinfectant, comprising the step of:
electrolytically generating silver in a solution of citric acid and water to formed an aqueous solution of silver citrate.

22. The process of making a disinfectant as set forth in claim 21, wherein the step of electrolytically generating silver includes forming an organic metal complex with the citric acid.
23. The process of making a disinfectant as set forth in claim 21, wherein the step of electrolytically generating silver includes forming a chelated organic metal complex with the citric acid.
24. The process of making a disinfectant as set forth in claim 21, wherein the step of electrolytically generating silver includes forming a complex with the citric acid of $(Ag(CA)_x)^+ (CA)^-$, wherein CA is $(C_6H_8O_7 - H_2O)$.
25. The process of making a disinfectant as set forth in claim 21, wherein the step of electrolytically generating silver includes forming a complex with the citric acid of $(Ag+CA)^-$, wherein CA is $(C_6H_8O_7 - H_2O)$.
26. The process of making an improved aqueous disinfectant, comprising the step of:
- creating a solution of approximately 5.0% to 10% citric acid in water by volume;
 - spacing a positive silver electrode relative to a negative electrode for enabling the solution to be located therebetween;
 - applying a potential difference to the positive and negative electrodes to establish a flow of silver ions between the positive and negative electrodes for enabling the silver ions to react with the citric acid to form silver citrate thereby.
27. The process of making an improved aqueous disinfectant as set forth in claim 26, wherein the step of spacing a positive silver electrode relative to a negative electrode includes spacing the positive silver electrode from the negative electrode a distance sufficient to enable silver ion flow therebetween.

28. The process of making an improved aqueous disinfectant as set forth in claim 26, wherein the step of spacing a positive silver electrode relative to a negative electrode includes spacing the positive silver electrode greater than 2.0 mm. from the negative electrode.

29. The process of making an improved aqueous disinfectant as set forth in claim 26, wherein the step of applying a potential difference to the positive and negative electrodes includes applying a potential difference to establish a flow of silver ions in the range of 0.1 amperes to 0.5 amperes.

30. An aqueous solution of silver citrate, comprising:
an aqueous solution of silver citrate in a solution of citric acid and water wherein the concentration of silver citrate exceeds 0.05% by volume.

31. The process of making silver citrate, comprising the step of:
electrolytically generating silver in a solution of citric acid and water to formed an aqueous solution of silver citrate.

32. The process of making silver citrate as set forth in claim 31, wherein the step of electrolytically generating silver includes forming an organic metal complex with the citric acid.

33. The process of making silver citrate as set forth in claim 31, wherein the step of electrolytically generating silver includes forming a chelated organic metal complex with the citric acid.

34. The process of making silver citrate as set forth in claim 31, wherein the step of electrolytically generating silver includes forming a complex with the citric acid of $(Ag(CA)_x)^+ (CA)^-$, wherein CA is $(C_6H_8O_7 - H_2O)$.

35. The process of making silver citrate as set forth in claim 31, wherein the step of electrolytically generating silver includes forming a complex with the citric acid of (Ag+CA⁻), wherein CA is (C₆H₈O₇ - H₂O).




FIG. 1




FIG. 2




FIG. 3




FIG. 4

INITIAL INTERVAL SHELF-LIFE TEST

	<u>1% Citric Acid</u>			<u>10% Citric Acid</u>		
	<u>3 ppm</u>	<u>30 ppm</u>	<u>300 ppm</u>	<u>3 ppm</u>	<u>30 ppm</u>	<u>300 ppm</u>
Week 1	1.006	9.630	2.060	0.676	0.690	145.94
	0.983	9.925	1.960	0.434	1.010	157.32
	1.174	9.765	1.490	0.688	0.865	165.34
Avg.	1.054	9.773	1.837	0.599	0.855	156.20
Week 2	0.877	0.455	<0.050	0.858	4.580	158.62
	0.649	0.310	<0.050	0.950	4.635	157.10
	0.738	0.290	<0.050	0.870	4.765	159.18
Avg.	0.755	0.352	<0.050	0.893	4.660	158.30
Week 3	0.630	0.925	2.690	1.140	15.060	160.62
	0.688	2.150	6.600	1.230	27.025	163.44
	0.666	3.380	0.110	1.370	26.605	158.68
Avg.	0.661	2.152	3.133	1.247	22.897	160.91
Week 4	0.624	2.115	1.260	3.094	3.355	151.82
	0.652	0.850	8.830	2.094	4.865	163.10
	0.600	1.055	0.140	2.582	7.975	169.96
Avg.	0.625	1.340	1.743	2.590	5.398	161.63

FIG. 5

SECONDARY INTERVAL SHELF-LIFE TEST

	<u>1.0% Citr Acid</u>	<u>5.0% citric Acid</u>	<u>10% Citric Acid</u>
Week 0	772.0 mg/L	282.0 mg/L	678.0 mg/L
Week 7	143.3 mg/L	359.2 mg/L	824.6 mg/L
Week 14	139.8 mg/L	359.6 mg/L	842.5 mg/L
	143.6 mg/L	355.9 mg/L	822.0 mg/L
	140.5 mg/L	358.2 mg/L	817.3 mg/L
Mean	141.3 mg/L	357.9 mg/L	827.3 mg/L
Week 21	1.063 mg/L	350.2 mg/L	982.9 mg/L
	1.022 mg/L	335.0 mg/L	1019.4 mg/L
	1.066 mg/L	331.0 mg/L	1017.3 mg/L
	1.050 MYLL	338.7 mg/L	1006.5 mg/L

FIG. 6

EFFICACY AGAINST SALMONELLA CHOLERASUIS

<u>Sample</u>	<u>15 sec.</u>	<u>1 Minute</u>	<u>5 Minutes</u>	<u>10 minutes</u>	<u>30 minutes</u>
Ethanol 20% (a)	1.93E+06	7.10E+05	5.20E+03	2.00E+02	1.00E+02
Ethanol 20% (b)	2.38E+06	1.02E+06	7.10E+03	3.00E+02	1.00E+02
Ethanol 20% (c)	1.66E+06	6.70E+05	8.90E+03	1.00E+02	1.00E+02
Ethanol 20% (d)	1.95E+06	6.70E+05	8.50E+03	1.00E+02	1.00E+02
Average	1.98E+06	7.68E+05	7.43E+03	1.75E+02	1.00E+02
% Reduction	81.3208	92.7594	99.9300	99.9983	99.9991
Ag Citric Acid & ETOH (a)	6.00E+03	1.00E+01	1.00E+01	1.00E+01	1.00E+01
Ag Citric Acid & ETOH (b)	6.80E+03	1.00E+01	1.00E+01	1.00E+01	1.00E+01
Ag Citric Acid & ETOH (c)	5.27E+03	1.00E+01	1.00E+01	1.00E+01	1.00E+01
Ag Citric Acid & ETOH (d)	4.81E+03	1.00E+01	1.00E+01	1.00E+01	1.00E+01
Average	5.72E+03	1.00E+01	1.00E+01	1.00E+01	1.00E+01
% Reduction	99.9460	99.9999	99.9999	99.9999	99.9999
Citric Acid (a)		1.35E+06	8.00E+01	1.00E+01	1.00E+01
Citric Acid (b)		1.22E+06	5.00E+01	1.00E+01	1.00E+01
Citric Acid (c)		1.00E+06	1.00E+02	1.00E+01	1.00E+01
Citric Acid (d)		1.27E+06	7.00E+01	1.00E+01	1.00E+01
Average		1.21E+06	7.50E+01	1.00E+01	1.00E+01
% Reduction		88.5849	99.9993	99.9999	99.9999
Inoculum (a)				7.30E+06	
Inoculum (b)				1.39E+07	
Average				1.06E+07	

FIG. 7

EFFICACY AGAINST STAPHYLOCOCCUS AUREUS

Sample	15 sec.	1 Minute	5 Minutes	10 minutes	30 minutes
Ethanol 20% (a)	4.29E+06	9.30E+06	6.80E+06	9.30E+06	7.20E+06
Ethanol 20% (b)	9.30E+06	8.40E+06	9.70E+06	8.40E+06	1.06E+07
Ethanol 20% (c)	1.06E+07	8.20E+06	7.40E+06	8.30E+06	8.50E+06
Ethanol 20% (d)	1.00E+07	8.20E+06	8.10E+06	8.50E+06	1.23E+07
Average	8.55E+06	8.53E+06	8.10E+06	8.63E+06	9.65E+06
% Reduction	37.8364	38.0000	41.8182	37.2727	29.8182
Ag Citric Acid & Ethanol (a)	2.54E+06	5.51E+04	1.00E+01	1.00E+01	1.00E+01
Ag Citric Acid & Ethanol (b)	2.67E+06	2.50E+04	1.00E+01	1.00E+01	1.00E+01
Ag Citric Acid & Ethanol (c)	6.83E+06	8.00E+04	1.00E+01	1.00E+01	1.00E+01
Ag Citric Acid & Ethanol (d)	4.46E+06	8.10E+04	1.00E+01	1.00E+01	1.00E+01
Average	4.13E+06	6.03E+04	1.00E+01	1.00E+01	1.00E+01
% Reduction	70.0000	99.5616	99.999	99.9999	99.9999
Ag Citric Acid (a)		6.99E+06	1.06E+06	9.10E+03	1.00E+01
Ag Citric Acid (b)		5.50E+06	6.90E+05	8.30E+03	1.00E+01
Ag Citric Acid (c)		7.31E+06	8.00E+05	1.89E+04	1.00E+01
Ag Citric Acid (d)		5.87E+06	9.70E+05	1.66E+04	1.00E+01
Average		6.42E+06	8.80E+05	1.32E+04	1.00E+01
% Reduction		53.3271	93.6000	99.9038	99.9999
Citric Acid (a)		7.10E+06	1.06E+07	1.17E+07	1.13E+07
Citric Acid (b)		8.20E+06	9.70E+06	1.18E+07	7.40E+06
Citric Acid (c)		1.05E+07	8.60E+06	6.70E+06	2.80E+06
Citric Acid (d)		6.00E+06	9.60E+06	7.30E+06	1.06E+07
Average		7.95E+06	9.63E+06	9.38E+06	8.03E+06
% Reduction		42.1818	30.0000	31.8182	41.6364
Inoculum (a)			1.41E+07		
Inoculum (b)			1.34E+07		
Average			1.38E+07		

FIG. 8

EFFICACY AGAINST PSEUDOMONAS AERUGINOSA

Sample	15 sec.	1 min.	5 mins.	10 mins.	30 mins.
Ethanol 20% (a)	2.85E+06	2.95E+06	8.10E+06	3.70E+06	2.30E+06
Ethanol 20% (b)	3.26E+06	2.90E+06	7.30E+06	3.90E+06	2.25E+06
Ethanol 20% (c)	2.37E+06	2.50E+06	6.70E+06	3.14E+06	5.70E+06
Ethanol 20% (d)	2.55E+06	2.50E+06	7.50E+06	3.00E+06	6.90E+06
Average	2.76E+06	2.71E+06	7.40E+06	3.44E+06	4.29E+06
% Reduction	66.9760	67.5150	11.3772	58.8623	48.6527
Ag citric Acid & ETOH (a)	9.60E+05	1.58E+04	1.00E+01	1.00E+01	1.00E+01
Ag Citric Acid & ETOH (b)	1.27E+06	1.43E+04	1.00E+01	1.00E+01	1.00E+01
Ag Citric Acid & ETOH (c)	1.58E+06	1.40E+04	1.00E+01	3.00E+01	1.00E+01
Ag Citric Acid & ETOH (d)	1.43E+04	1.41E+04	1.00E+01	5.00E+01	1.00E+01
Average	9.56E+05	1.46E+04	1.00E+01	2.50E+01	1.00E+01
% Reduction	88.5500	99.8257	99.9999	99.9997	99.9999
Ag Citric Acid (a)		3.17E+06	9.10E+05	1.51E+04	1.00E+01
Ag Citric Acid (b)		2.55E+06	1.38E+06	2.22E+04	1.00E+01
Ag Citric Acid (c)		3.08E+06	1.13E+06	3.91E+04	Lost count
Ag Citric Acid (d)		2.64E+06	1.03E+06	3.17E+04	3.00E+01
Average		2.86E+06	1.11E+06	2.70E+04	1.67E+01
% Reduction		65.7485	86.6766	99.6763	99.9998
Citric Acid (a)		7.90E+06	3.03E+06	7.30E+06	7.50E+06
Citric Acid (b)		6.90E+06	2.70E+06	7.70E+06	2.90E+06
Citric Acid (c)		1.29E+07	2.88E+07	2.22E+06	6.10E+06
Citric Acid (d)		1.00E+07	2.72E+07	7.90E+06	2.80E+06
Average		9.43E+06	1.54E+07	6.28E+06	4.83E+06
% Reduction		-12.8743	-84.8204	24.7904	42.2156
Inoculum (a)			7.80E+06		
Inoculum (b)			8.90E+06		
Average			8.35E+06		

FIG. 9

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US98/21604
A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A01N 37/04, 55/02, 59/00; A61K 31/19, 31/28, 33/38

US CL : 424/618, 619; 422/22, 28; 514/495, 574, 724

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/618, 619; 422/22, 28; 514/495, 574, 724

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	SRIVASTAVA, G.C. et al. Development of ready to use antiseptic dressings - Part I: development of surgical gauze and absorbent cotton by using in vitro laboratory evaluation methods. Labdev J. Sci Tech. October 1970, Vol. 8-B, No. 4, pages 209-213, especially pages 209 and Tables 1 and 3 at pages 211-212.	1-8, 10 ----- 1-20
X --- Y	Chem. abstr., Vol. 87, No. 10, 05 September 1977 (Columbus, OH, USA), page 403, column 2, the abstract No. 74283n, TSIMBLER, S.M. et al. 'Complexes of silver(I) with some hydroxy acids.' Zh. Neorg. Khim. 1977, 22(7), 1842-1846 (Russ.).	1-8, 10 ----- 1-10

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:		*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A	document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B	earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A*	document member of the same patent family
O	document referring to an oral disclosure, use, exhibition or other means		
P	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

19 JANUARY 1999

Date of mailing of the international search report

02 FEB 1999

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JOHN PAK

Telephone No. 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/21604

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 4,055,655 A (MAURER et al.) 25 October 1977, column 1, lines 59-65, column 2, lines 31-38, column 4, lines 1-9, column 5, lines 15-35, column 13, lines 36-37, claims 1-3 and 8.	1-10
A	JP 47-32537 B (FUKUDA Y.) 19 August 1972, see the attached English abstract.	1-35

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/21604

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

•
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/21604

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

STN ONLINE, Files CAPLUS, WPIDS, AGRICOLA, CABA, BIOSIS. Search terms: silver, Ag, citric, citrate#,
propanetricarboxyl?, electrol?, microampere?, ampere?, amps, current#, anode#, cathode#, volt?, electrode#.